arXiv:1401.7964v2 [math.GN] 1 Feb 2014

METRIZABLE DH-SPACES OF THE FIRST CATEGORY

SERGEY MEDVEDEV

ABSTRACT. We show that if a separable space X has a meager open subset containing a copy of the Cantor set 2^ω, then X has $\mathfrak{c}$ types of countable dense subsets. We suggest a generalization of the λ-set for non-separable spaces. Let X be an h-homogeneous Λ-set. Then X is densely homogeneous and $X \setminus A$ is homeomorphic to X for every σ-discrete subset $A \subset X$.

All spaces under discussion are metrizable.

We see the rapid growth of the theory of CDH-spaces lately. For example, K. Kunen, A. Medini and L. Zdomskyy [KMZ] proved that if a separable metrizable space X is not CB but has a CB dense subset, then X has $\mathfrak{c}$ types of countable dense subsets. They obtained a similar result [KMZ, Theorem 16] for a non-Baire space with the perfect set property for open sets. In fact, it suffices (see Theorem 1.4) to have a meager open subset containing a copy of the Cantor set 2^ω.

We introduce the Λ-sets as a generalization of the λ-sets for non-separable spaces and consider its properties. In particular, we improve (see Theorem 2.2) the result due to R. Hernández-Gutiérrez, M. Hrušák and J. van Mill [HHM, Proposition 4.9] concerning CDH-property of h-homogeneous λ-sets.

In the paper we are not dealing with the set-theoretic methods; only topological methods are applied.

1. DH-spaces

For all undefined terms and notation see [En].

$X \approx Y$ means that X and Y are homeomorphic spaces. A separable topological space X is *countable dense homogeneous* (briefly, CDH) if, given any two countable dense subsets A and B of X, there is a homeomorphism $h : X \to X$ such that $h(A) = B$. The *type* of a countable dense subset D of a separable space X is $\{h(D) : h \text{ is a homeomorphism of } X\}$. So a separable space is CDH if and only if it has exactly one type of countable dense subsets. Also notice that the maximum possible number of types of countable dense subsets of a separable space is $\mathfrak{c}$.

A space X is *densely homogeneous* (briefly, DH) provided that (1) X has a σ-discrete dense subset and (2) if A and B are two σ-discrete dense subsets of X, then there is a homeomorphism $h : X \to X$ such that $f(A) = B$. One can check that no DH-space without isolated points can be a σ-discrete space. Clearly, if X is a separable metrizable space, then X is CDH $\Leftrightarrow$ X is DH.

2010 *Mathematics Subject Classification.* 54H05, 54E52.

Key words and phrases. h-homogeneous space, set of first category, dense homogeneous, countable dense homogeneous, Λ-space.

A space X is called *of the first category* (or *meager*) if it can be represented as a countable union of nowhere dense sets. A space X is *completely Baire* (briefly, CB) if every closed subspace of X is a Baire space.

Lemma 1.1 was obtained by the author [M2]. Independently it was proved for separable spaces by B. Fitzpatrick Jr. and H-X. Zhou [FZ].

Lemma 1.1. *For a metric space X the following are equivalent:*

1) *the space X is of the first category,*

2) *X contains a σ-discrete everywhere dense set of type G_δ without isolated points.*

Recall that a separable space in which every countable set is a G_δ-set is called a *λ-set*. Likewise, a space in which every σ-discrete set is a G_δ-set is called a *Λ-set*. From Lemma 1.1 it follows that every metrizable Λ-set without isolated points is of the first category in itself.

The following statement is similar to [FZ, Theorem 3.4].

Theorem 1.2. *Every DH, meager, metrizable space X is a Λ-set.*

Proof. Take a σ-discrete subset A of X. By Lemma 1.1, there is a σ-discrete dense G_δ-set $B \subset X$. Clearly, $A \cup B$ is a σ-discrete dense subset of X. Then $A \cup B$ is a G_δ-set in X because $A \cup B = h(B)$ for some homeomorphism $h : X \to X$. Since A is a G_δ-set in $A \cup B$, A is a G_δ-set in X. □

Theorem 1.3. *If a meager metrizable space X contains a copy of the Cantor set 2^ω, then X is not DH.*

Proof. Let $2^\omega \approx F \subset X$. Clearly, F is a closed nowhere dense subset of X. Take a dense set $A \subset F$ such that A is homeomorphic to the rationals. By Lemma 1.1, there is a σ-discrete dense G_δ-set $B \subset X$. The set $F \setminus (A \cup B)$ is not F_σ in F because it is homeomorphic to the irrationals. Hence, $X \setminus (A \cup B)$ is not F_σ in X. Since $A \cup B$ is not G_δ in X, there is no homeomorphism $f : X \to X$ with $f(B) = A \cup B$. Thus, X is not DH. □

Theorem 1.4. *Suppose a separable metrizable space X has a meager open subset containing a copy of the Cantor set 2^ω. Then X has $\mathfrak{c}$ types of countable dense subsets.*

Proof. The set $V = \bigcup\{U : U \text{ is a meager open set in } X\}$ is the largest open subset of X which is meager in itself. One can check that $Y = \{y \in X : \text{every neighborhood of } y \text{ contains a copy of the Cantor set } 2^\omega\}$ is a closed subset of X. Under the conditions of the theorem, $V \cap Y \neq \emptyset$.

Case 1. Let $V \cap \mathrm{Int}(Y) = \emptyset$. Then $V \cap Y$ is a non-empty closed nowhere dense subset of V. Choose a countable set $D \subset V$ such that $V \subseteq \overline{D}$ and $D \cap Y = \emptyset$. If $X \setminus V \neq \emptyset$, fix a countable set D^* such that $\overline{D^*} = X \setminus V$; otherwise put $D^* = \emptyset$. Take a copy $K \subset V \cap Y$ of the Cantor set 2^ω and consider a family $\{E_\alpha \subset K : \alpha < \mathfrak{c}\}$ of size $\mathfrak{c}$ consisting of pairwise non-homeomorphic countable spaces; such family exists by the theorem due to S. Mazurkiewicz and W. Sierpiński

[MS]. For every $\alpha < \mathfrak{c}$, define $D_\alpha = E_\alpha \cup D \cup D^*$. Clearly, each D_α is a countable dense subset of X.

We claim that D_α and D_β are countable dense subsets of X of a different type whenever $\alpha \neq \beta$. Assume that there exists a homeomorphism $h : X \to X$ such that $h(D_\alpha) = D_\beta$. One readily sees that $h(V) = V$, $h(X \setminus V) = X \setminus V$, and $h(V \cap Y) = V \cap Y$. Hence, $E_\beta = D_\beta \cap V \cap Y = h(D_\alpha \cap V \cap Y) = h(E_\alpha)$. This contradicts the fact that E_α and E_β are non-homeomorphic.

Case 2. Let $V^* = V \cap \mathrm{Int}(Y) \neq \emptyset$. Now we will use ideas from the proof of [HM, Theorem 4.5]. Fix a copy $K \subset V^*$ of the Cantor set 2^ω. Clearly, K is nowhere dense in V^*. Then there exists an open set W such that $\overline{W} \subset V^*$, $V^* \setminus \overline{W} \neq \emptyset$, and the boundary $\mathrm{Fr}(W) = K$. Obviously, $W \cap K = \emptyset$. Using Lemma 1.1, we can choose a countable dense subset $D \subset W$ which is a G_δ-set in X. Take a countable base $\{V_n : n \in \omega\}$ for $V^* \setminus \overline{W}$. For every $n \in \omega$ we may pick a closed nowhere dense set $F_n \subset V_n$ such that $F_n \approx 2^\omega$. Without loss of generality, $F_n \cap F_m = \emptyset$ whenever $n \neq m$. Let D_n be a countable dense subset of F_n. Clearly, each D_n is homeomorphic to the rationals. Put $D' = \cup\{D_n : n \in \omega\}$. Note that $D' \cap O$ is not a G_δ-subset of X for every non-empty open set $O \subseteq V^* \setminus \overline{W}$. For if this were true, we could pick $n \in \omega$ such that $F_n \subset O$ which would imply that $D' \cap F_n = D_n$ would be G_δ in F_n. The last contradicts to the Baire category theorem because the rationals cannot be homeomorphic to a G_δ-subset of a compact space.

Fix a countable dense subset D'' of $X \setminus \overline{V^*}$. Put $D^* = D \cup D' \cup D''$.

As above, there exists a family $\{E_\alpha \subset K : \alpha < \mathfrak{c}\}$ of size $\mathfrak{c}$ consisting of pairwise non-homeomorphic countable spaces. For every $\alpha < \mathfrak{c}$, define $D_\alpha = E_\alpha \cup D^*$. Clearly, each D_α is a countable dense subset of X.

We claim that D_α and D_β are countable dense subsets of X of a different type whenever $\alpha \neq \beta$. Assume that there exists a homeomorphism $h : X \to X$ such that $h(D_\alpha) = D_\beta$. One readily sees that $h(V) = V$ and $h(Y) = Y$. Hence, $h(V^*) = V^*$. Moreover, let us show that $h(W) \cap (V^* \setminus \overline{W}) = \emptyset$. Assume the converse. Then there is a non-empty open set $O \subset W$ such that $h(O) \subset V^* \setminus \overline{W}$. However, this is impossible because $O \cap D$ is a G_δ-set in X while $h(O \cap D) = h(O) \cap D'$ is not. Likewise, we obtain $W \cap h(V^* \setminus \overline{W}) = \emptyset$. Thus, $h(W) = W$ and $h(V^* \setminus \overline{W}) = V^* \setminus \overline{W}$. Hence, $h(K) = K$ because $V^* = W \cup K \cup (V^* \setminus \overline{W})$. Then $E_\beta = D_\beta \cap K = h(D_\alpha \cap K) = h(E_\alpha)$. This contradicts the fact that E_α and E_β are non-homeomorphic. □

2. h-Homogeneous DH-spaces

A space X is called *h-homogeneous* if $\mathrm{Ind}X = 0$ and every non-empty clopen subset of X is homeomorphic to X. If $\mathcal{U}$ is a family of subsets of a metric space (X, ϱ), then $\cup\mathcal{U} = \cup\{U : U \in \mathcal{U}\}$ and $\mathrm{mesh}(\mathcal{U}) = \sup\{\mathrm{diam}U : U \in \mathcal{U}\}$ is a *measure* of $\mathcal{U}$.

Lemma 2.1. *Let X_i be a metrizable space with $\mathrm{Ind}X_i = 0$ and F_i be a nowhere dense closed subset of X_i, where $i \in \{1,2\}$. Let $f : F_1 \to F_2$ be a homeomorphism.*

Then there exist a σ-discrete (in X_i) cover $\mathcal{V}_i$ of $X_i \setminus F_i$ by non-empty pairwise disjoint clopen subsets of X_i, $i \in \{1,2\}$, and a bijection $\psi : \mathcal{V}_1 \to \mathcal{V}_2$ such that for any subsets $D_1 \subseteq \bigcup \mathcal{V}_1$ and $D_2 \subseteq \bigcup \mathcal{V}_2$ and any bijection $g : D_1 \to D_2$ satisfying $g(D_1 \cap V) = D_2 \cap \psi(V)$ for every $V \in \mathcal{V}_1$, the combination mapping $f \triangledown g : F_1 \cup D_1 \to F_2 \cup D_2$ is continuous at each point of F_1 and its inverse $(f \triangledown g)^{-1}$ is continuous at each point of F_2.

Under the conditions of Lemma 2.1, we say [M4] that:
1) the cover $\mathcal{V}_i$ *forms a residual family* with respect to F_i, where $i \in \{1,2\}$,
2) the bijection $\psi : \mathcal{V}_1 \to \mathcal{V}_2$ is *agreed to* f.

Remark. A construction similar to Lemma 2.1 was used by B. Knaster and M. Reichbach [KR] for separable spaces and by A. V. Ostrovsky [Os] and the author [M3] for non-separable spaces. The last two papers were written in Russian, therefore the proof of Lemma 2.1 was also given in [M4]. Note that like statements are obtained for arbitrary metrizable spaces in [M3] and [M4].

Independently Lemma 2.1 was obtained by F. van Engelen [vE]. In his notation, the triple $\langle \mathcal{V}_1, \mathcal{V}_2, \psi \rangle$ is called a *Knaster-Reichbach cover*, or *KR-cover*, for $\langle X_1 \setminus F_1, X_2 \setminus F_2, f \rangle$. Sometimes this term is more convenient.

Theorem 2.2. *Let X be an h-homogeneous metrizable Λ-set. Then X is* **DH**.

Proof. Let $w(X) = \tau$. Take two σ-discrete dense subsets A and B of X. From h-homogeneity of X it follows that A and B are weight-homogeneous spaces of weight τ. Then $A \approx B \approx Q(\tau)$ by [M1, Theorem 1], where $Q(\tau)$ is the small σ-product of countably many discrete spaces of cardinality τ. Let $A = \cup\{A_n : n \in \omega\}$, where each A_n is discrete in X, and $B = \cup\{B_n : n \in \omega\}$, where each B_n is discrete in X.

Fix a metric ϱ on X which induces the original topology on X. Since $\mathrm{Ind}X = \dim X = 0$, by [En, Theorem 7.3.1], there exists a sequence $\mathcal{U}_0, \mathcal{U}_1, \ldots$ of discrete clopen covers of X such that $\mathrm{mesh}(\mathcal{U}_n) < 1/(n+1)$ and $\mathcal{U}_{n+1}$ is a refinement of $\mathcal{U}_n$ for each n. Obviously, the family $\mathcal{U} = \{U \in \mathcal{U}_n : n \in \omega\}$ forms a base for X. For each $U \in \mathcal{U}$ fix a homeomorphism $\varphi_U : X \to U$. Then the sets $A^* = A \cup \bigcup\{\varphi_U(A) : U \in \mathcal{U}\}$ and $B^* = B \cup \bigcup\{\varphi_U(B) : U \in \mathcal{U}\}$ are σ-discrete in X. By definition of a Λ-set, A^* and B^* are both G_δ-subsets of X. Since $X^* = X \setminus (A^* \cup B^*)$ is a dense subset of X, it is a meager subset of X. Then X^* can be represented as $\cup\{X_n : n \in \omega\}$, where each X_n is a nowhere dense closed subset of X. By construction, $\varphi_U(X_n) \cap A = \emptyset$ and $\varphi_U(X_n) \cap B = \emptyset$ for any $n \in \omega$ and $U \in \mathcal{U}$.

The difference $X \setminus (A \cup X^*)$ is a σ-discrete subset of X. Then $X \setminus (A \cup X^*) = \cup\{A_n^* : n \in \omega\}$, where each non-empty A_n^* is discrete in X. Similarly, $X \setminus (B \cup X^*) = \cup\{B_n^* : n \in \omega\}$, where each non-empty B_n^* is discrete in X.

We will construct the homeomorphism $f : X \to X$ satisfying $f(A) = B$ by induction on n. For each $n \in \omega$ we will find a pair of closed nowhere dense sets F_n and E_n, a homeomorphism $f_n : F_n \to E_n$, a KR-cover $\langle \mathcal{V}_n, \mathcal{W}_n, \psi_n \rangle$ for $\langle X \setminus F_n, X \setminus E_n, f_n \rangle$ such that:

(a) $F_n \cup X_n \cup A_n^* \cup A_n \subset F_{n+1}$,

(b) $E_n \cup X_n \cup B_n^* \cup B_n \subset E_{n+1}$,

(c) $f_n(A \cap F_n) = B \cap E_n$,

(d) the restriction $f_n|_{F_i} = f_i$ for every $i < n$,

(e) $\mathrm{mesh}(\mathcal{V}_n) < 1/(n+1)$ and $\mathrm{mesh}(\mathcal{W}_n) < 1/(n+1)$,

(f) $\mathcal{V}_{n+1}$ refines $\mathcal{V}_n$ and $\mathcal{W}_{n+1}$ refines $\mathcal{W}_n$,

(g) if $V \in \mathcal{V}_n$ and $V^* \in \mathcal{V}_m$ for $n < m$, then $V^* \subset V$ if and only if $\psi_m(V^*) \subset \psi_n(V)$.

For $n = 0$, let $F_0 = E_0 = \emptyset$, $\mathcal{V}_0 = \mathcal{W}_0 = \{X\}$, $\psi_0(X) = X$, and $f_0(\emptyset) = \emptyset$.

Now let us consider the induction step $n + 1$ for some $n \in \omega$.

First, fix $V \in \mathcal{V}_n$ and $W = \psi_n(V) \in \mathcal{W}_n$. Let $Y = V \cap X_m$, where m is the least i with $V \cap X_i \neq \emptyset$, and $Z = W \cap X_l$, where l is the least i with $W \cap X_i \neq \emptyset$.

Since A is dense in X and X is nowhere locally compact and weight-homogeneous, there exists a discrete (in X) set $A' \subset A \cap V$ of cardinality τ such that $A_m \cap V \subset A'$, where m is the least i with $V \cap A_i \neq \emptyset$. We can also find a discrete (in X) set $B' \subset B \cap W$ of cardinality τ such that $B_l \cap W \subset B'$, where l is the least i with $W \cap B_i \neq \emptyset$. Choose a bijection $g : A' \to B'$.

Next, let $A^{*\prime} = V \cap A_m^*$, where m is the least i with $V \cap A_i^* \neq \emptyset$, and $B^{*\prime} = W \cap B_l^*$, where l is the least i with $W \cap B_i^* \neq \emptyset$. Now we have to distinguish some possibilities. Case 1: $A^{*\prime} \neq \emptyset$ and $B^{*\prime} \neq \emptyset$. By adding points from $V \cap (X^* \setminus Y)$ or from $W \cap (X^* \setminus Z)$, if it is necessary, we may assume that $A^{*\prime}$ and $B^{*\prime}$ are both discrete subsets of X of the same cardinality. Case 2: $A^{*\prime} \neq \emptyset$ and $B^{*\prime} = \emptyset$. In this case we replace $B^{*\prime}$ by a discrete subset from $W \cap (X^* \setminus Z)$ of cardinality $|A^{*\prime}|$. Case 3: $A^{*\prime} = \emptyset$ and $B^{*\prime} \neq \emptyset$ is similar to Case 2. Case 4: $A^{*\prime} = B^{*\prime} = \emptyset$. For Cases 1–3 we can take a bijection $h : A^{*\prime} \to B^{*\prime}$. For Case 4 we put $h(\emptyset) = \emptyset$.

The union $Y' = Y \cup A' \cup A^{*\prime}$ is a closed nowhere dense subset of V. Hence there exists a basic set $U \in \mathcal{U}$ such that $U \subset V$ and $U \cap Y' = \emptyset$. Then $\varphi_U(Z) \approx Z$ and $\varphi_U(Z)$ is a closed nowhere dense subset of V. Define $F_V = Y' \cup \varphi_U(Z)$. Likewise, there exists $U^* \in \mathcal{U}$ such that $U^* \subset W$ and $\varphi_{U^*}(Y) \approx Y$ misses the set $Z' = Z \cup B' \cup B^{*\prime}$. Put $E_W = Z' \cup \varphi_{U^*}(Y)$.

Then we define a homeomorphism $f_V : F_V \to E_W$ by the rule

$$f_V(x) = \begin{cases} g(x) & \text{if } x \in A', \\ h(x) & \text{if } x \in A^{*\prime}, \\ \varphi_{U^*}(x) & \text{if } x \in Y, \\ (\varphi_U)^{-1}(x) & \text{if } x \in \varphi_U(Z). \end{cases}$$

By Lemma 2.1, there exists a KR-cover $\langle \mathcal{V}_V, \mathcal{W}_V, \psi_V \rangle$ for $\langle V \setminus F_V, W \setminus E_W, f_V \rangle$.

It is not hard to see that $F_{n+1} = F_n \cup \bigcup\{F_V : V \in \mathcal{V}_n\}$ and $E_{n+1} = E_n \cup \bigcup\{E_{\psi_n(V)} : V \in \mathcal{V}_n\}$ are closed nowhere dense subsets of X. Clearly, the mapping

$$f_{n+1}(x) = \begin{cases} f_V(x) & \text{if } x \in F_V \text{ for some } V \in \mathcal{V}_n, \\ f_n(x) & \text{if } x \in F_n \end{cases}$$

is a homeomorphism between F_{n+1} and E_{n+1}.

Define $\mathcal{V}_{n+1} = \{U \in \mathcal{V}_V : V \in \mathcal{V}_n\}$ and $\mathcal{W}_{n+1} = \{\psi_V(U) : U \in \mathcal{V}_V, V \in \mathcal{V}_n\}$. Then $\psi_{n+1} = \bigcup\{\psi_V : V \in \mathcal{V}_n\}$ is a bijection between $\mathcal{V}_{n+1}$ and $\mathcal{W}_{n+1}$ which is agreed to f_{n+1}. Thus, $\langle \mathcal{V}_{n+1}, \mathcal{W}_{n+1}, \psi_{n+1}\rangle$ is a KR-cover for $\langle X \setminus F_{n+1}, X \setminus E_{n+1}, f_{n+1}\rangle$. If necessary, the families $\mathcal{V}_{n+1}$ and $\mathcal{W}_{n+1}$ can be refined to families with measure less than $1/(n+2)$. One can check that all conditions (a)–(g) are satisfied. This completes the induction step.

From (a) and (b) it follows that $X = \bigcup\{F_n : n \in \omega\} = \bigcup\{E_n : n \in \omega\}$. The condition (d) implies that the rule $f(x) = f_n(x)$ if $x \in F_n$ for some n defines the bijection $f : X \to X$. Moreover, f is a homeomorphism. Next, using (c), we obtain $f(A) = B$. □

Corollary 2.3. *Let $X \subset 2^\omega$ be an h-homogeneous λ-set. Then X is* CDH.

Theorem 2.4. *Let X be an h-homogeneous metrizable Λ-set. If A is a σ-discrete subset of X, then X is homeomorphic to $X \setminus A$.*

Proof. Since $\dim X = 0$, there exists a sequence $\mathcal{U}_0, \mathcal{U}_1, \dots$ of discrete clopen covers of X such that $\mathcal{U}_{n+1}$ is a refinement of $\mathcal{U}_n$ for each n and the family $\mathcal{U} = \{U \in \mathcal{U}_n : n \in \omega\}$ forms a base for X. For each $U \in \mathcal{U}$ fix a homeomorphism $\varphi_U : X \to U$. Then the set $A^* = A \cup \bigcup\{\varphi_U(A) : U \in \mathcal{U}\}$ is σ-discrete in X. Hence, $A^* = \cup\{A_n : n \in \omega\}$, where each A_n is a discrete closed subset of X. For every $U \in \mathcal{U}$ the set $U \setminus A$ contains a closed copy of each A_n because the space X is weight-homogeneous and nowhere locally compact.

By definition of a Λ-set, A^* is a G_δ-subset of X. Since $X^* = X \setminus A^*$ is a dense subset of X, it is meager in X. Then $X^* = \cup\{X_n : n \in \omega\}$, where each X_n is a nowhere dense closed subset of X. By construction, $\varphi_U(X_n)$ is a closed nowhere dense subset of U and $\varphi_U(X_n) \subset U \setminus A$ for any $n \in \omega$ and $U \in \mathcal{U}$.

Since $X = \bigcup\{A_n \cup X_n : n \in \omega\}$, by [M4, Lemma 4], every non-empty relatively open subset of $X \setminus A$ contains a closed (in $X \setminus A$) nowhere dense copy of X. According to [M4, Theorem 3], $X \setminus A$ is homeomorphic to $h(X, k)$, where $k = w(X)$ and $h(X, k)$ is the smallest h-homogeneous meager space containing X. Clearly, $X = h(X, k)$. □

Corollary 2.5. *Let $X \subset 2^\omega$ be an h-homogeneous λ-set. If A is a countable subset of X, then X is homeomorphic to $X \setminus A$.*

References

[FZ] B. Fitzpatrick Jr. and H-X. Zhou, Countable dense homogeneity and the Baire property, Topology Applic., 43 (1992) 1–14.

[vE] F. van Engelen, On the homogeneity of infinite products, Topol. Proc., 17 (1992) 303–315.

[En] R. Engelking, General topology, PWN, Warszawa, 1977.

[HHM] R. Hernández-Gutiérrez, M. Hrušák and J. van Mill, Countable dense homogeneity and λ-sets, preprint, 2013.

[HM] M. Hrušák and J. van Mill, Nearly countable dense homogeneous spaces, preprint, 2012, to appear in *Canad. J. Math.*.

[KR] B. Knaster and M. Reichbach, Notion d'homogénéité et prolongements des homéomorphies, Fund. Math., 40 (1953) 180–193.

[KMZ] K. Kunen, A. Medini and L. Zdomskyy, Seven characterizations of non-meager P-filters, 2013, arxiv 1311.1677.

[MS] S. Mazurkiewicz and W. Sierpiński, Contribution à la topologie des ansembles dénombrables, Fund. Math., 1:1 (1920) 17–27.

[M1] S. V. Medvedev, Topological characteristics of the space $Q(k)$ and $Q \times B(k)$, Vestn. Mosk. Univ., Ser. I, Mat. Mekh. 41:1 (1986) 47–49 (in Russian). English transl.: Mosc. Univ. Math. Bull. 41:1 (1986) 42–45.

[M2] S. V. Medvedev, On a problem for spaces of the first category, Vestn. Mosk. Univ., Ser. I, Mat. Mekh. 41:2 (1986) 84–86 (in Russian). English transl.: Mosc. Univ. Math. Bull. 41:2 (1986) 62–65.

[M3] S. V. Medvedev, On the structure of metric h-homogeneous spaces, in: V. V. Fedorčuk, A. V. Arhangel'skii, V. I. Zaĭcev (Eds.), General topology. Mappings of topological spaces, Moscow State Univ. 1986, pp. 77–98 (in Russian).

[M4] S. V. Medvedev, About closed subsets of spaces of first category, Topology Appl., 159:8 (2012) 2187–2192.

[Os] A. V. Ostrovsky, Continuous images of the product $C \times Q$ of the Cantor perfect set C and the rational numbers Q, in: Seminar on General Topology, Moskov. Gos. Univ., Moscow, 1981, 78–85 (in Russian).

South Ural State University, Chelyabinsk, 454080 Russia
E-mail address: medv@math.susu.ac.ru